Subject: [radiojove-data] 17 Feb 2016 lo-B/D, non-lo-A

From: Dave Typinski <davetyp@typnet.net>

Date: 02/20/2016 12:10

To: RadioJove-Data <radiojove-data@lists.nasa.gov>

Adding to Wes & Jim & Tom & John & Nathan's nice reports, here's some lo-B/D and non-lo-A from Wednesday morning.

RCP dominant L bursting 0425-0714 UTC from 16 to 27 MHz, vertex early arcs (lo-B).

RCP dominant S bursting 0427-0437 UTC from 19 to 23 MHz, zero drift emission envelope (lo-B).

LCP dominant narrow band S burst trains 0529-0557 UTC from 18 to 21 MHz, positive drift emission envelopes (first lo-D arc).

LCP dominant L bursting 0708-0807 UTC from 16 to 21 MHz, positive drift emission envelope (second lo-D arc).

RCP dominant L bursting 0649-0819 UTC from 16 to 26 MHz, negative drift emission envelope (non-lo-A).

It is known that Io-D sometimes produces two arcs in series; in this case the conditions were such that the second arc (the LCP L bursting) was easier to observe than the first arc (the LCP narrow band S burst trains).


Jupiter was -42° to +17° off axis.


Jupiter was leading the Sun by 157°.


Dave


AJ4CO Observatory 17 Feb 2016, log entry 606 - 610


1 of 6














6 of 6